

FEVERAL[®]

IBUPROFEN ORAL SUSPENSION 100 mg/5 mL

1-INDICATIONS:

Feveral[®] contains ibuprofen which belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs) indicated to relieve pain and fever due to colds or flu, sore throat, headache, earache, toothache and immunization. It provides up to 8 hours relief of fever and aches.

2-COMPOSITION:

Each 5 mL of Feveral[®] contains 100 mg of ibuprofen, carboxymethylcellulose sodium, citric acid, EDTA sodium, glycerin, sodium benzoate, polysorbate 80, propylene glycol, sorbitol solution, sucralose, acesulfame potassium, xanthan gum, strawberry flavor, red 40, microcrystalline cellulose, purified water.

3-MECHANISM OF ACTION:

Ibuprofen, a non-steroidal anti-inflammatory drug (NSAID), is a non selective inhibitor of cyclooxygenase enzymes COX-1 and COX-2 involved in prostaglandin (mediators of pain and fever) synthesis via the arachidonic acid pathway.

4-DOSAGE AND ADMINISTRATION:

To be taken orally.

Shake well before use.

Doses below may be repeated every 6-8 hours if needed. Do not use more than 4 times a day.

Age (Years)	Weight	Dosage
2-3	10.9-15.9 kg	1¼ teaspoons = 6 mL
4-5	16.0-21.3 kg	2 teaspoons = 10 mL
6-8	21.4-26.7 kg	2½ teaspoons = 12.5 mL
9-10	26.8-32.5 kg	3 teaspoons = 15 mL
11-12	32.6-43.0 kg	3¾ teaspoons = 19 mL

Children below 2 years: Please ask your doctor.

Children over 12 years of age and adults who have difficulty swallowing tablets may take 200 mg every 4 hours (2 teaspoons) or 400 mg every 6-8 hours (4 teaspoons). Do not exceed 1200 mg a day (12 teaspoons).

5- SIDE EFFECTS:

If your child notices any of the following side effects, stop giving this medicine and seek immediate medical help:

- Allergic reaction: skin reddening, facial swelling, hives, blisters
- Wheezing
- Stomach bleeding: Stop giving immediately this medicine and talk to a doctor if your child vomits blood, has black or bloody stools and stomach pain, The risk is higher if the child has a history of bleeding problems and stomach ulcers or if he takes anticoagulants or other drugs containing NSAIDs.
- Sore throat with high fever, headache, nausea and vomiting.
- High blood pressure.
- Heart attack and stroke: Chest pain, trouble breathing, weakness in one part of the body, slurred speech and leg swelling. The risk is higher if this medicine is given more than directed or for a longer time than directed.

6- WARNINGS AND PRECAUTIONS:

Ask a doctor before use If your child:

- Is taking any other drug.
- Is under a doctor's care for any serious condition.
- Has stomach ulcers and bleeding problems.
- Has ever had an allergic reaction to any other pain reliever.
- Has high blood pressure or heart problems.
- Has severe kidney or liver failure.
- Has severe dehydration.

Do not use **Feveral**[®] before or after heart surgery.

Stop giving **Feveral**[®] to your child and ask a doctor if fever or pain gets worse or lasts more than 3 days.

Do not give your child more than the directed dose.

7- STORAGE:

- Store below 25°C.
- Protect from light.
- Keep out of reach of children.

8- PACKAGING AND VOLUME:

DESCRIPTION	CODE	SHELF LIFE
Feveral [®] 100 mL	30 01100	2 years

Manufactured by:



Tel: +961 9 235 490

Fax: 09 230 489 - 09 235 489

www.alfalabs.com.lb

©ALFALABSSAL |  Alfa Labs S.A.L.