

PerioKIN mouthwash

0.20% CHLORHEXIDINE DG

0.10% CETYLPYRIDINIUM CHLORIDE

High concentration of chlorhexidine for an efficacious and complete oral protection.

- Colourless: transparent solution for an easier identification of blood when rinsing.
- Oral hygiene during dental interventions.

Indications

- Oral hygiene in cases of:
- Localized gingival disorders.
- Post-surgery.
- After tooth extractions.
- Scaling and root planing.
- Curettage.
- Implant procedures.
- Patients with physical disabilities.
- Patients with temporomandibular joint disorders.



PerioKIN gel

0.20% CHLORHEXIDINE DG

Chlorhexidine efficacy targeted exactly where needed

High bioadhesiveness


PerioKIN Gel formulation confers great adherence and bioadhesiveness, which allows it to act on the affected area.

Indications

- Oral hygiene in cases of:
- Localized gingival disorders.
- Post-surgery.
- After tooth extractions.
- Scaling and root planing.
- Curettage.
- Implant procedures.



Maximum adhesion and bioadhesivity



Representative model of molecular interpenetration of the bioadhesive polymer A, with the glycoproteins of the mucosa B. (a), (b) and (c) represent the consecutive stages of approaching, superficial contact and mucoadhesive polymer/mucus glycoprotein interpenetration.